

Lie group analysis of Poisson's equation and optimal system of subalgebras for Lie algebra of 3-dimensional rigid motions

M. Nadjafikhah ^{a,★}

^a*School of Mathematics, Iran University of Science and Technology, Narmak-16, Tehran, I.R.IRAN.*

Abstract

Using the basic Lie symmetry method, we find the most general Lie point symmetries group of the $\nabla u = f(u)$ Poisson's equation, which has a subalgebra isomorphic to the 3-dimensional special Euclidean group $SE(3)$ or group of rigid motions of $\mathbb{R}^3$. Looking the adjoint representation of $SE(3)$ on its Lie algebra $\mathfrak{se}(3)$, we will find the complete optimal system of its subalgebras. This latter provides some properties of solutions for the Poisson's equation.

Key words: Lie-point symmetries, similarity solution, adjoint representation, optimal system for Lie subalgebras.

1 Introduction

The *Poisson's equation* is

$$\nabla u = f(u), \tag{1}$$

where $\nabla = \partial^2/\partial x^2 + \partial^2/\partial y^2 + \partial^2/\partial z^2$, $f : \mathbb{R} \rightarrow \mathbb{R}$ is a given smooth function, Ω is an open subset of $\mathbb{R}^3$ and $u : \Omega \rightarrow \mathbb{R}$ is an unknown smooth function [2]. This is a stationary heat equation with a nonlinear source (5.4.1 of [6]). In this paper, using the basic Lie symmetry method, we find the most general Lie point symmetries group of the Poisson's equation, which has a subalgebra isomorphic to 3-dimensional special Euclidean group $SE(3)$. Looking the adjoint representation of $SE(3)$ on its Lie algebra $\mathfrak{se}(3)$, we will find the complete classification of its subalgebras up to conjugacy. This latter provide a process for building new solutions for the Poisson's equation.

Inspired by Galois' theory of, Sophus Lie developed an analogous theory of symmetry for differential equations. Lie's theory led to an algorithmic way to find special explicit solutions to differential equations with symmetry. These special solutions are called *group invariant solutions* and they constitute practically every known explicit solution to the systems of non-linear partial differential equations arising in mathematical physics, differential geometry and other areas.

These group-invariant solutions are found by solving a reduced system of differential equations involving fewer independent variables than the original system. For example, the solutions to a partial differential equation in two independent variables which are invariant under a given one-parameter symmetry group are all found by solving a system of ordinary differential equations. Today the search for group invariant solutions is still a common approach to explicitly solving non-linear partial differential equations. A good modern introduction to the use of similarity methods in engineering applications is the book [7]. An excellent introductory reference to Lie's theory of group invariant solutions is [4].

★ Corresponding author: Tel. +98-21-73913426. Fax +98-21-77240472.
Email address: m_nadjafikhah@iust.ac.ir (M. Nadjafikhah).

2 Lie point symmetries

In this section, we find the Lie point symmetries of Poisson's equation, using the basic prolongation method and the infinitesimal criterion of invariance as [4]. Consider the one parameter Lie group of infinitesimal transformations on $(x_1 = x, x_2 = y, x_3 = z, u)$ given by

$$\tilde{x}_i = x^i + s \xi_i(x_1, x_2, x_3, u) + O(s^2), \quad \tilde{u} = u + s \varphi(x_1, x_2, x_3, u) + O(s^2), \quad i = 1, 2, 3, \quad (2)$$

where s is the group parameter and ξ_i , $i = 1, 2, 3$ and φ are the infinitesimals of the transformations for the independent and dependent variables, respectively. The associated vector field is $\mathbf{v} = \sum_{i=1}^3 \xi_i \partial_{x_i} + \varphi \partial_u$, and its second prolongation is

$$\text{Pro}^{(2)} \mathbf{v} = \mathbf{v} + \varphi^x \partial_{u_x} + \varphi^y \partial_{u_y} + \varphi^z \partial_{u_z} + \varphi^{xx} \partial_{u_{xx}} + \varphi^{xy} \partial_{u_{xy}} + \varphi^{xz} \partial_{u_{xz}} + \varphi^{yy} \partial_{u_{yy}} + \varphi^{yz} \partial_{u_{yz}} + \varphi^{zz} \partial_{u_{zz}}, \quad (3)$$

with

$$\begin{aligned} \varphi^x &= -u_x(\partial_x + u_x \partial_u) \xi_1 - u_y(\partial_x + u_x \partial_u) \xi_2 - u_z(\partial_x + u_x \partial_u) \xi_3 + (\partial_x + u_x \partial_u) \varphi, \\ &\vdots \\ \varphi^{zz} &= -(2u_{xz} \partial_z + (u_x u_{zz} + 2u_z u_{xz}) \partial_u - u_x \partial_{zz} - 2u_x u_z \partial_{zu} - u_x u_z^2 \partial_{uu}) \xi_1 \\ &\quad - (2u_{yz} \partial_z + u_y u_{zz} + 2u_z u_{yz}) \partial_u + u_y \partial_{zz} + 2u_y u_z \partial_{zu} + u_y u_z^2 \partial_{uu}) \xi_2 \\ &\quad - (2u_{zz} \partial_z + 3u_{zz} u_z \partial_u + u_z \partial_{zz} + 2u_z^2 \partial_{zu} + u_z^3 \partial_{uu}) \xi_3 + (u_{zz} \partial_u + \partial_{zz} + 2u_z \partial_{zu} + u_z^2 \partial_{uu}) \varphi. \end{aligned} \quad (4)$$

The vector field $\mathbf{v}$ generates a one parameter symmetry group of Poisson's equation if and only if

$$\text{Pro}^{(2)} \mathbf{v}(\nabla u - f(u)) = 0 \quad \text{whenever} \quad \nabla u = f(u). \quad (5)$$

This condition is equivalent to the set of defining equations:

$$\begin{aligned} \partial_u \xi_1 &= 0, & \partial_x \xi_1 &= \partial_z \xi_3, \\ \partial_u \xi_2 &= 0, & \partial_x \xi_2 + \partial_y \xi_1 &= 0, & \partial_{zz} \xi_2 + \partial_{yy} \xi_2 + \partial_{xx} \xi_2 &= 2\partial_{yu} \varphi, \\ \partial_u \xi_3 &= 0, & \partial_x \xi_3 + \partial_z \xi_1 &= 0, & \partial_{zz} \xi_3 + \partial_{yy} \xi_3 + \partial_{xx} \xi_3 &= 2\partial_{zu} \varphi, & \nabla \varphi &= 2F(u) \cdot \partial_z \xi_3 + F'(u) \cdot \varphi. \\ \partial_{uu} \varphi &= 0, & \partial_y \xi_3 + \partial_z \xi_2 &= 0, & \partial_{xx} \xi_1 + \partial_{zz} \xi_1 + \partial_{yy} \xi_1 &= 2\partial_{xu} \varphi, \\ & & \partial_z \xi_3 &= \partial_y \xi_2, \end{aligned} \quad (6)$$

By solving this system of PDEs, we find that

2.0.0.1 Theorem *The Lie group $\mathbf{g}_f$ of point symmetries of Poisson's equation (1) is an infinite dimensional Lie algebra generated by the vector fields $\mathbf{X} = \xi_1 \partial_x + \xi_2 \partial_y + \xi_3 \partial_z + \varphi \partial_u$, where*

$$\begin{aligned} \xi_1 &= a_7(x^2 - y^2 - z^2) + 2(a_5 y + a_1 z)x + a_6 x + a_8 y - a_4 z + a_9, \\ \xi_2 &= a_5(y^2 - z^2 - x^2) + 2(a_7 x + a_1 z)y - a_8 x + a_6 y + a_2 z + a_{10}, & \varphi &= F_1(x, y, z) \cdot u + F_2(x, y, z), \\ \xi_3 &= a_1(z^2 - x^2 - y^2) + 2(a_7 x + a_5 y)z + a_4 x - a_2 y + a_6 z + a_3, \end{aligned} \quad (7)$$

with

$$F_1 = a_{11} - (a_7 x + a_5 y + a_1 z), \quad \nabla F_2 = (5F_1 - 2a_6 - 4a_{11}) \cdot f(u) - (uF_1 + F_2) \cdot f'(u), \quad (8)$$

and a_i , $i = 1, \dots, 11$ are arbitrary constants.

Table 1

Commutation table of $\mathfrak{g}_f$.

	X_1	X_2	X_3	X_4	X_5	X_6
X_1	0	0	0	0	$-X_3$	X_2
X_2	0	0	0	X_3	0	$-X_1$
X_3	0	0	0	$-X_2$	X_1	0
X_4	0	$-X_3$	X_2	0	$-X_6$	X_5
X_5	X_3	0	$-X_1$	X_6	0	$-X_4$
X_6	$-X_2$	X_1	0	$-X_5$	X_4	0

2.0.0.2 Conclusion The Lie group $\mathfrak{g}_f$ of point symmetries of Poisson's equation (1) has a 6-dimensional Lie subalgebra generated by the following vector fields:

$$X_1 = \partial_x, \quad X_2 = \partial_y, \quad X_3 = \partial_z, \quad X_4 = y\partial_z - z\partial_y, \quad X_5 = z\partial_x - x\partial_z, \quad X_6 = x\partial_y - y\partial_x. \quad (9)$$

The commutator table of $\mathfrak{g}_f$ is given in Table 1, where the entry in the i^{th} row and j^{th} column is defined as $[X_i, X_j] = X_i \cdot X_j - X_j \cdot X_i$, $i, j = 1, \dots, 6$. $\square$

An isometry (or rigid motion) of Euclidean space $\mathbb{R}^3$ is a mapping $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ which preserves the Euclidean distance between points. For every isometry F , there exist a unique transformation $T \in \text{Trans}(3) \cong \mathbb{R}^3$ and a unique orthogonal transformation $C \in \text{O}(3)$ such that $F = C \circ T$. $\text{Trans}(3)$ is a normal subgroup of $\text{E}(3)$, and $\text{E}(3) \cong \mathbb{R}^3 \rtimes \text{O}(3)$. $\text{E}(3)$ has a natural 6-dimensional Lie group structure. The connected component containing I_3 is *special Euclidean group* $\text{SE}(3)$, consists of all orientation preserving isometries, and is a connected 6-dimensional Lie group. Furthermore, the Lie algebra $\mathfrak{se}(3)$ of 3-dimensional special Euclidean group is spanned by the following 6 vector fields (9).

2.0.0.3 Remark The $\mathfrak{g}_f$ Lie group of point symmetries of Poisson's equation $\nabla u = f(u)$ may be changed, for special choices of f ; for example, if $f = 0$, that is for $\nabla u = 0$ Laplace equation, $\mathfrak{g}_f$ is spanned by X_i , $i = 1, \dots, 6$, ∂_u , $\partial_x + \partial_y + \partial_z$, $xz\partial_x + yz\partial_y + (z^2 - x^2 - y^2)\partial_z + zu\partial_u$, $(x^2 - y^2 - z^2)\partial_x + xy\partial_y + xz\partial_z + xu\partial_u$, $xy\partial_x + (x^2 + y^2 + z^2)\partial_y + yz\partial_z + yu\partial_u$, and so on.

3 Optimal system of subalgebras

Let a system of differential equation Δ admitting the symmetry Lie group G , be given. Now G operates on the set of solutions S of Δ . Let $s \cdot G$ be the orbit of s , and H be a subgroup of G . Invariant H -solutions $s \in S$ are characterized by equality $s \cdot S = \{s\}$. If $h \in G$ is a transformation and $s \in S$, then $h \cdot (s \cdot H) = (h \cdot s) \cdot (hHh^{-1})$. Consequently, every invariant H -solution s transforms into an invariant hHh^{-1} -solution (Proposition 3.6 of [4]).

Therefore, different invariant solutions are found from similar subgroups of G . Thus, classification of invariant H -solutions is reduced to the problem of classification of subgroups of G , up to similarity. An optimal system of s -dimensional subgroups of G is a list of conjugacy inequivalent s -dimensional subgroups of G with the property that any other subgroup is conjugate to precisely one subgroup in the list. Similarly, a list of s -dimensional subalgebras forms an optimal system if every s -dimensional subalgebra of $\mathfrak{g}$ is equivalent to a unique member of the list under some element of the adjoint representation: $\tilde{\mathfrak{h}} = \text{Ad}(g) \cdot \mathfrak{h}$.

Let H and $\tilde{H}$ be connected, s -dimensional Lie subgroups of the Lie group G with corresponding Lie subalgebras $\mathfrak{h}$ and $\tilde{\mathfrak{h}}$ of the Lie algebra $\mathfrak{g}$ of G . Then $\tilde{H} = gHg^{-1}$ are conjugate subgroups if and only if $\tilde{\mathfrak{h}} = \text{Ad}(g) \cdot \mathfrak{h}$ are conjugate subalgebras (Proposition 3.7 of [4]). Thus, the problem of finding an optimal system of subgroups is equivalent to that of finding an optimal system of subalgebras, and so we concentrate on it.

For one-dimensional subalgebras, the classification problem is essentially the same as the problem of classifying the orbits of the adjoint representation, since each one-dimensional subalgebra is determined by a nonzero vector in $\mathfrak{se}(3)$ and so to "simplify" it as much as possible.

The adjoint action is given by the Lie series

$$\text{Ad}(\exp(s.X_i).X_j) = X_j - s.[X_i, X_j] + \frac{s^2}{2}.[X_i, [X_i, X_j]] - \dots, \quad (10)$$

where $[X_i, X_j]$ is the commutator for the Lie algebra, s is a parameter, and $i, j = 1, \dots, 6$ ([4], page 199). We can write the adjoint action for the Lie algebra $\mathfrak{se}(3)$, and show that

3.0.0.4 Theorem *A one-dimensional optimal system of $\mathfrak{se}(3)$ is given by*

$$\begin{aligned} A_1^1 : X_6, & \quad A_1^3 : X_2 + b X_5, & A_1^5 : X_1 + a X_2 + b X_6, & \quad A_1^7 : X_2 + a X_3 + b X_5, \\ A_1^2 : X_1 + b X_4, & A_1^4 : X_3 + b X_6, & A_1^6 : X_1 + a X_3 + b X_4, \end{aligned} \quad (11)$$

where $a, b \in \mathbb{R}$ and $a \neq 0$.

Proof: $F_i^s : \mathfrak{se}(3) \rightarrow \mathfrak{se}(3)$ defined by $X \mapsto \text{Ad}(\exp(s_i X_i).X)$ is a linear map, for $i = 1, \dots, 6$. The matrix M_i^s of F_i^s , $i = 1, \dots, 6$, with respect to basis $\{X_1, \dots, X_6\}$ is

$$\begin{aligned} M_1^s &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & s & 0 & 1 & 0 \\ 0 & -s & 0 & 0 & 0 & 1 \end{bmatrix}, & M_2^s &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -s & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ s & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, & M_3^s &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & s & 0 & 1 & 0 & 0 \\ -s & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \\ M_4^s &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & C & S & 0 & 0 & 0 \\ 0 & S & C & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & C & S \\ 0 & 0 & 0 & 0 & -S & C \end{bmatrix}, & M_5^s &= \begin{bmatrix} C & 0 & -S & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ S & 0 & C & 0 & 0 & 0 \\ 0 & 0 & 0 & C & 0 & -S \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & S & 0 & C \end{bmatrix}, & M_6^s &= \begin{bmatrix} C & S & 0 & 0 & 0 & 0 \\ -S & C & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & C & S & 0 \\ 0 & 0 & 0 & -S & C & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}. \end{aligned} \quad (12)$$

respectively, where $S = \sin s$ and $C = \cos s$. Let $X = \sum_{i=1}^6 a_i X_i$, then

$$F_6^{s_6} \circ \dots \circ F_1^{s_1} : X \mapsto (\cos s_5 \cdot \cos s_6 \cdot a_1 + \cos s_5 \cdot \sin s_6 \cdot a_2 - \sin s_5 \cdot a_3).X_1 + \dots \quad (13)$$

Now, we can simply X as follows:

If a_1, a_2 and $a_3 = 0$, then we can make the coefficients of $X_1, \dots, X_5$ vanish, by $s_4 = -\arctan(a_5/a_6)$ and $s_5 = \arctan(a_4/a_6)$. Scaling X if necessary, we can assume that $a_6 = 1$. And X is reduced to the Case of A_1^1 .

If a_2 and $a_3 = 0$ but $a_1 \neq 0$, then we can make the coefficients of X_2, X_3, X_5 and X_6 vanish, by $s_2 = -a_6/a_1$ and $s_3 = a_5/a_1$. Scaling X if necessary, we can assume that $a_1 = 1$. And X is reduced to the Case of A_1^2 .

If a_1 and $a_3 = 0$ but $a_2 \neq 0$, then we can make the coefficients of X_1, X_2, X_3 , and X_6 vanish, by $s_1 = a_6/a_2$ and $s_3 = -a_4/a_2$. Scaling X if necessary, we can assume that $a_3 = 1$. And X is reduced to the Case of A_1^3 .

If a_1 and $a_2 = 0$ but $a_3 \neq 0$, then we can make the coefficients of X_1, X_2, X_4 and X_5 vanish, by $s_1 = -a_5/a_3$ and $s_2 = a_4/a_3$. Scaling X if necessary, we can assume that $a_3 = 1$. And X is reduced to the Case of A_1^4 .

If a_1 and $a_2 \neq 0$, then we can make the coefficients of X_3 , X_4 and X_5 vanish, by $s_1 = a_6/a_2$, $s_3 = a_5/a_1$ and $s_4 = -\arctan(a_3/a_2)$. Scaling X if necessary, we can assume that $a_1 = 1$. And X is reduced to the Case of A_1^5 .

If $a_2 = 0$ but a_1 and $a_3 \neq 0$, then we can make the coefficients of X_2 , X_5 and X_6 vanish, by $s_1 = -a_5/a_3$, $s_2 = -a_6/a_1$ and $s_4 = -\arctan(a_3/a_2)$. Scaling X if necessary, we can assume that $a_1 = 1$. And X is reduced to the Case of A_1^6 .

If $a_1 = 0$ but a_2 and $a_3 \neq 0$, then we can make the coefficients of X_3 , X_4 and X_5 vanish, by $s_1 = a_6/a_2$, and $s_2 = a_4/a_3$. Scaling X if necessary, we can assume that $a_2 = 1$. And X is reduced to the Case of A_1^7 . $\square$

3.0.0.5 Theorem A two-dimensional optimal system of $\mathfrak{se}(3)$ is given by

$$\begin{aligned} A_2^1 : X_2, X_5, & \quad A_2^3 : X_1, X_2 + a X_4, & \quad A_2^5 : X_2, X_3 + a X_5, \\ A_2^2 : X_3, X_6, & \quad A_2^4 : X_1, X_3 + a X_4, & \quad A_2^6 : X_3, X_1 + a X_4, \end{aligned} \quad (14)$$

where $a \in \mathbb{R}$. All of these sub-algebras are Abelian.

Proof: In this proof, we shall assume that $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X, Y\}$ is a 2-dimensional Lie sub-algebra of $\mathfrak{se}(3)$.

Let $X = X_6$ be as the Case of A_1^1 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_3 X_3 + b_6 X_6$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $Y = X_3$. Now $\mathfrak{g}$ is reduced to the Case of A_2^2 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$ defined as (12).

Let $X = X_1 + a_6 X_6$ be as the Case of A_1^2 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_3 X_3 + b_1 X$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $Y = X_3$. Now $\mathfrak{g}$ is reduced to the Case of A_2^6 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$.

Let $X = X_2 + a_5 X_5$ be as the Case of A_1^3 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_2 X_2 + b_5 X_5$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_2$ and $Y = X_5$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^1 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$.

Let $X = X_3 + a_6 X_6$ be as the Case of A_1^4 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_3 X_3 + b_6 X_6$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_6$ and $Y = X_3$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^2 .

Let $X = X_1 + a_2 X_2 + a_4 X_4$ be as the Case of A_1^5 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_1 X_1 + (b_2/a_2) X$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_1$ and $Y = X_2 + a X_4$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^3 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$.

Let $X = X_2 + a_3 X_3 + a_4 X_4$ be as the Case of A_1^6 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_1 X_1 + (b_3/a_3) X$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_1$ and $Y = X_3 + a X_4$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^4 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$.

Let $X = X_2 + a_3 X_3 + a_5 X_5$ be as the Case of A_1^7 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_2 X_2 + (b_3/a_3) X$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_2$ and $Y = X_3 + a X_5$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^5 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$. And, we have proved the Theorem. $\square$

3.0.0.6 Theorem A three-dimensional optimal system of $\mathfrak{se}(3)$ is given by $A_3 : X_1 + a X_4, X_2, X_3$, where $a \in \mathbb{R}$. The commutator table of A_3 is Table 2.

Table 2
Commutation table of A_3 .

	X	Y	Z
X	0	$-a$	0
Y	a	0	0
Z	0	0	0

Table 3
Commutation table of A_4 .

	X_1	X_2	X_3	X_4
X_1	0	0	0	0
X_2	0	0	0	X_3
X_3	0	0	0	$-X_2$
X_4	0	$-X_3$	X_2	0

Proof: In this proof, we assume that $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X, Y, Z\}$ is a 3-dimensional Lie sub-algebra of $\mathfrak{se}(3)$.

Let $X = X_2$ and $Y = X_5$ are as the Case of A_2^1 , $Z = \sum_{i=1}^6 b_i X_i$, $[X, Z] = \alpha_1 X + \beta_2 Y + \beta_3 Z$ and $[Y, Z] = \beta_1 X + \beta_2 Y + \beta_3 Z$. Then, we have $Z = b_2 X + b_5 Y$. By a suitable change of base of $\mathfrak{g}$, we can assume that $Z = 0$, and $\mathfrak{g}$ is not a 3-dimensional sub-algebra. Thus, in this case we have not any 3-dimensional sub-algebra.

The Cases A_2^i , $i = 2, 3, 4, 5$ are similar. Thus, in these cases we have not any 3-dimensional sub-algebra.

Let $X = X_3$ and $Y = X_1 + a X_4$ are as the Case of A_2^6 , $Z = \sum_{i=1}^6 b_i X_i$, $[X, Z] = \alpha_1 X + \beta_2 Y + \beta_3 Z$ and $[Y, Z] = \beta_1 X + \beta_2 Y + \beta_3 Z$. Then, we have $Z = (b_1/\alpha_3) \cdot (-\beta_3 X + \alpha_3 Y - a X_2)$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_1 + a X_4$, $Y = X_2$ and $Z = X_3$. Now, $\mathfrak{g}$ is reduced to the Case of A_3 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$ defined as (12). $\square$

3.0.0.7 Theorem A four-dimensional optimal system of $\mathfrak{se}(3)$ is given by $A_4 : X_1, X_2, X_3, X_4$. The commutator table of A_4 is Table 3. *Proof:* Assume that $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X_1 + a X_4, X_2, X_3, X\}$ be a Lie sub-algebra of $\mathfrak{se}(3)$, where

$X = \sum_{i=1}^6 b_i X_i$. Then, we have $X = \sum_{i=1}^4 b_i X_i$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_4$. Now, $\mathfrak{g}$ is reduced to the Case of A_4 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$ defined as (12). $\square$

3.0.0.8 Theorem $\mathfrak{se}(3)$ has not any five-dimensional Lie sub-algebra.

Proof: Assume that $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X_1, X_2, X_3, X_4, X\}$ be a Lie sub-algebra of $\mathfrak{se}(3)$, where $X = \sum_{i=1}^6 b_i X_i$. Then, we have $X = \sum_{i=1}^4 b_i X_i$, and $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X_1, X_2, X_3, X_4\}$ is not a 5-dimensional sub-algebra. $\square$

4 Invariant solutions

To obtain the group transformation which is generated by the infinitesimal generators $X_i = \sum_{j=1}^3 \xi_j^i + \phi^i \partial_u$ for $i = 1, \dots, 6$, we need to solve the 6 systems of first order ordinary differential equations,

$$\begin{aligned} \tilde{x}'_j(s) &= \xi_j^i(\tilde{x}_1(s), \tilde{x}_2(s), \tilde{x}_3(s), \tilde{u}(s)), \quad \tilde{x}_j(0) = x_j, \quad i = 1, \dots, 6 \\ \tilde{u}'(s) &= \phi^i(\tilde{x}_1(s), \tilde{x}_2(s), \tilde{x}_3(s), \tilde{u}(s)), \quad \tilde{u}(0) = u. \quad j = 1, 2, 3 \end{aligned} \tag{15}$$

Exponentiating the infinitesimal symmetries of the Poisson's equation (1), we get the one parameter groups $g_k(s)$ generated by X_k for $k = 1, \dots, 6$; and consequently, we have

4.0.0.9 Theorem If $u = h(x, y, z)$ is a solution of the Poisson's equation (1), so are the functions

$$\begin{aligned} g_1(s) \cdot h(x, y, z) &= h(x + s, y, z), & g_4(s) \cdot h(x, y, z) &= h(x, y \cos s - z \sin s, z \cos s + y \sin s), \\ g_2(s) \cdot h(x, y, z) &= h(x, y + s, z), & g_5(s) \cdot h(x, y, z) &= h(x \cos s + z \sin s, y, z \cos s - x \sin s), \\ g_3(s) \cdot h(x, y, z) &= h(x, y, z + s), & g_6(s) \cdot h(x, y, z) &= h(x \cos s - y \sin s, x \sin s + y \cos s, z), \end{aligned} \quad (16)$$

References

- [1] G.W. BLUMAN, J.D. COLE, *Similarity Methods for Differential Equations*, Springer, Berlin, 1974.
- [2] J. JOST, *Partial differential equations*, Springer, GTM 214, 2002.
- [3] M. NADJAFIKHAH, *Lie Symmetries of Inviscid Burgers Equation*, Adv. appl. Clifford alg., DOI 10.1007/s00006-003-0000.
- [4] P.J. OLVER, *Applications of Lie Groups to Differential Equations*, Springer, New York, 1986.
- [5] L.V. OVSIANNIKOV, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.
- [6] A.D. POLYANIN, and V.F. ZAITSEV, *Handbook of Nonlinear Partial Differential Equations*, Chapman & Hall/CRC, Boca Raton, 2004.
- [7] R. SESHADRI and T.Y. NA, *Group Invariance in Engineering Boundary Value Problems*, Springer-Verlag, New York, 1985.
- [8] T.R. SEKHAR, V.D. SHARMA, *Similarity solutions for three dimensional Euler equations using Lie group analysis*, Appl. Math. and Comp. 196 (2008) 147–157.

Lie group analysis of Poisson's equation and optimal system of subalgebras for Lie algebra of 3-dimensional rigid motions

M. Nadjafikhah ^{a,★}

^a*School of Mathematics, Iran University of Science and Technology, Narmak, Tehran, Iran.*

Abstract

Using the basic Lie symmetry method, we find the most general Lie point symmetries group of the $\nabla u = f(u)$ Poisson's equation, which has a subalgebra isomorphic to the 3-dimensional special Euclidean group $SE(3)$ or group of rigid motions of $\mathbb{R}^3$. Looking the adjoint representation of $SE(3)$ on its Lie algebra $\mathfrak{se}(3)$, we will find the complete optimal system of its subalgebras. This latter provides some properties of solutions for the Poisson's equation.

Key words: Lie-point symmetries, similarity solution, adjoint representation, optimal system for Lie subalgebras.

1 Introduction

The *Poisson's equation* is

$$\nabla u = f(u), \tag{1}$$

where $\nabla = \partial^2/\partial x^2 + \partial^2/\partial y^2 + \partial^2/\partial z^2$, $f : \mathbb{R} \rightarrow \mathbb{R}$ is a given smooth function, Ω is an open subset of $\mathbb{R}^3$ and $u : \Omega \rightarrow \mathbb{R}$ is an unknown smooth function [2]. This is a stationary heat equation with a nonlinear source (5.4.1 of [6]). In this paper, using the basic Lie symmetry method, we find the most general Lie point symmetries group of the Poisson's equation, which has a subalgebra isomorphic to 3-dimensional special Euclidean group $SE(3)$. Looking the adjoint representation of $SE(3)$ on its Lie algebra $\mathfrak{se}(3)$, we will find the complete classification of its subalgebras up to conjugacy. This latter provide a process for building new solutions for the Poisson's equation.

Inspired by Galois' theory of, Sophus Lie developed an analogous theory of symmetry for differential equations. Lie's theory led to an algorithmic way to find special explicit solutions to differential equations with symmetry. These special solutions are called *group invariant solutions* and they constitute practically every known explicit solution to the systems of non-linear partial differential equations arising in mathematical physics, differential geometry and other areas.

These group-invariant solutions are found by solving a reduced system of differential equations involving fewer independent variables than the original system. For example, the solutions to a partial differential equation in two independent variables which are invariant under a given one-parameter symmetry group are all found by solving a system of ordinary differential equations. Today the search for group invariant solutions is still a common approach to explicitly solving non-linear partial differential equations. A good modern introduction to the use of similarity methods in engineering applications is the book [7]. An excellent introductory reference to Lie's theory of group invariant solutions is [4].

★ Corresponding author: Tel. +98-21-73913426. Fax +98-21-77240472.

Email address: m.nadjafikhah@iust.ac.ir (M. Nadjafikhah).

2 Lie point symmetries

In this section, we find the Lie point symmetries of Poisson's equation, using the basic prolongation method and the infinitesimal criterion of invariance as [4]. Consider the one parameter Lie group of infinitesimal transformations on $(x_1 = x, x_2 = y, x_3 = z, u)$ given by

$$\tilde{x}_i = x^i + s \xi_i(x_1, x_2, x_3, u) + O(s^2), \quad \tilde{u} = u + s \varphi(x_1, x_2, x_3, u) + O(s^2), \quad i = 1, 2, 3, \quad (2)$$

where s is the group parameter and ξ_i , $i = 1, 2, 3$ and φ are the infinitesimals of the transformations for the independent and dependent variables, respectively. The associated vector field is $\mathbf{v} = \sum_{i=1}^3 \xi_i \partial_{x_i} + \varphi \partial_u$, and its second prolongation is

$$\text{Pro}^{(2)} \mathbf{v} = \mathbf{v} + \varphi^x \partial_{u_x} + \varphi^y \partial_{u_y} + \varphi^z \partial_{u_z} + \varphi^{xx} \partial_{u_{xx}} + \varphi^{xy} \partial_{u_{xy}} + \varphi^{xz} \partial_{u_{xz}} + \varphi^{yy} \partial_{u_{yy}} + \varphi^{yz} \partial_{u_{yz}} + \varphi^{zz} \partial_{u_{zz}}, \quad (3)$$

with

$$\begin{aligned} \varphi^x &= -u_x(\partial_x + u_x \partial_u) \xi_1 - u_y(\partial_x + u_x \partial_u) \xi_2 - u_z(\partial_x + u_x \partial_u) \xi_3 + (\partial_x + u_x \partial_u) \varphi, \\ &\vdots \\ \varphi^{zz} &= -(2u_{xz} \partial_z + (u_x u_{zz} + 2u_z u_{xz}) \partial_u - u_x \partial_{zz} - 2u_x u_z \partial_{zu} - u_x u_z^2 \partial_{uu}) \xi_1 \\ &\quad - (2u_{yz} \partial_z + u_y u_{zz} + 2u_z u_{yz}) \partial_u + u_y \partial_{zz} + 2u_y u_z \partial_{zu} + u_y u_z^2 \partial_{uu}) \xi_2 \\ &\quad - (2u_{zz} \partial_z + 3u_{zz} u_z \partial_u + u_z \partial_{zz} + 2u_z^2 \partial_{zu} + u_z^3 \partial_{uu}) \xi_3 + (u_{zz} \partial_u + \partial_{zz} + 2u_z \partial_{zu} + u_z^2 \partial_{uu}) \varphi. \end{aligned} \quad (4)$$

The vector field $\mathbf{v}$ generates a one parameter symmetry group of Poisson's equation if and only if

$$\text{Pro}^{(2)} \mathbf{v}(\nabla u - f(u)) = 0 \quad \text{whenever} \quad \nabla u = f(u). \quad (5)$$

This condition is equivalent to the set of defining equations:

$$\begin{aligned} \partial_u \xi_1 &= 0, & \partial_x \xi_1 &= \partial_z \xi_3, \\ \partial_u \xi_2 &= 0, & \partial_x \xi_2 + \partial_y \xi_1 &= 0, & \partial_{zz} \xi_2 + \partial_{yy} \xi_2 + \partial_{xx} \xi_2 &= 2\partial_{yu} \varphi, \\ \partial_u \xi_3 &= 0, & \partial_x \xi_3 + \partial_z \xi_1 &= 0, & \partial_{zz} \xi_3 + \partial_{yy} \xi_3 + \partial_{xx} \xi_3 &= 2\partial_{zu} \varphi, & \nabla \varphi &= 2F(u) \cdot \partial_z \xi_3 + F'(u) \cdot \varphi. \\ \partial_{uu} \varphi &= 0, & \partial_y \xi_3 + \partial_z \xi_2 &= 0, & \partial_{xx} \xi_1 + \partial_{zz} \xi_1 + \partial_{yy} \xi_1 &= 2\partial_{xu} \varphi, \\ & & \partial_z \xi_3 &= \partial_y \xi_2, \end{aligned} \quad (6)$$

By solving this system of PDEs, we find that

2.0.0.1 Theorem *The Lie group $\mathbf{g}_f$ of point symmetries of Poisson's equation (1) is an infinite dimensional Lie algebra generated by the vector fields $\mathbf{X} = \xi_1 \partial_x + \xi_2 \partial_y + \xi_3 \partial_z + \varphi \partial_u$, where*

$$\begin{aligned} \xi_1 &= a_7(x^2 - y^2 - z^2) + 2(a_5 y + a_1 z)x + a_6 x + a_8 y - a_4 z + a_9, \\ \xi_2 &= a_5(y^2 - z^2 - x^2) + 2(a_7 x + a_1 z)y - a_8 x + a_6 y + a_2 z + a_{10}, & \varphi &= F_1(x, y, z) \cdot u + F_2(x, y, z), \\ \xi_3 &= a_1(z^2 - x^2 - y^2) + 2(a_7 x + a_5 y)z + a_4 x - a_2 y + a_6 z + a_3, \end{aligned} \quad (7)$$

with

$$F_1 = a_{11} - (a_7 x + a_5 y + a_1 z), \quad \nabla F_2 = (5F_1 - 2a_6 - 4a_{11}) \cdot f(u) - (uF_1 + F_2) \cdot f'(u), \quad (8)$$

and a_i , $i = 1, \dots, 11$ are arbitrary constants.

Table 1

Commutation table of $\mathfrak{g}_f$.

	X_1	X_2	X_3	X_4	X_5	X_6
X_1	0	0	0	0	$-X_3$	X_2
X_2	0	0	0	X_3	0	$-X_1$
X_3	0	0	0	$-X_2$	X_1	0
X_4	0	$-X_3$	X_2	0	$-X_6$	X_5
X_5	X_3	0	$-X_1$	X_6	0	$-X_4$
X_6	$-X_2$	X_1	0	$-X_5$	X_4	0

2.0.0.2 Conclusion The Lie group $\mathfrak{g}_f$ of point symmetries of Poisson's equation (1) has a 6-dimensional Lie subalgebra generated by the following vector fields:

$$X_1 = \partial_x, \quad X_2 = \partial_y, \quad X_3 = \partial_z, \quad X_4 = y\partial_z - z\partial_y, \quad X_5 = z\partial_x - x\partial_z, \quad X_6 = x\partial_y - y\partial_x. \quad (9)$$

The commutator table of $\mathfrak{g}_f$ is given in Table 1, where the entry in the i^{th} row and j^{th} column is defined as $[X_i, X_j] = X_i \cdot X_j - X_j \cdot X_i$, $i, j = 1, \dots, 6$. $\square$

An isometry (or rigid motion) of Euclidean space $\mathbb{R}^3$ is a mapping $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ which preserves the Euclidean distance between points. For every isometry F , there exist a unique transformation $T \in \text{Trans}(3) \cong \mathbb{R}^3$ and a unique orthogonal transformation $C \in \text{O}(3)$ such that $F = C \circ T$. $\text{Trans}(3)$ is a normal subgroup of $\text{E}(3)$, and $\text{E}(3) \cong \mathbb{R}^3 \rtimes \text{O}(3)$. $\text{E}(3)$ has a natural 6-dimensional Lie group structure. The connected component containing I_3 is *special Euclidean group* $\text{SE}(3)$, consists of all orientation preserving isometries, and is a connected 6-dimensional Lie group. Furthermore, the Lie algebra $\mathfrak{se}(3)$ of 3-dimensional special Euclidean group is spanned by the following 6 vector fields (9).

2.0.0.3 Remark The $\mathfrak{g}_f$ Lie group of point symmetries of Poisson's equation $\nabla u = f(u)$ may be changed, for special choices of f ; for example, if $f = 0$, that is for $\nabla u = 0$ Laplace equation, $\mathfrak{g}_f$ is spanned by X_i , $i = 1, \dots, 6$, ∂_u , $\partial_x + \partial_y + \partial_z$, $xz\partial_x + yz\partial_y + (z^2 - x^2 - y^2)\partial_z + zu\partial_u$, $(x^2 - y^2 - z^2)\partial_x + xy\partial_y + xz\partial_z + xu\partial_u$, $xy\partial_x + (x^2 + y^2 + z^2)\partial_y + yz\partial_z + yu\partial_u$, and so on.

3 Optimal system of subalgebras

Let a system of differential equation Δ admitting the symmetry Lie group G , be given. Now G operates on the set of solutions S of Δ . Let $s \cdot G$ be the orbit of s , and H be a subgroup of G . Invariant H -solutions $s \in S$ are characterized by equality $s \cdot S = \{s\}$. If $h \in G$ is a transformation and $s \in S$, then $h \cdot (s \cdot H) = (h \cdot s) \cdot (hHh^{-1})$. Consequently, every invariant H -solution s transforms into an invariant hHh^{-1} -solution (Proposition 3.6 of [4]).

Therefore, different invariant solutions are found from similar subgroups of G . Thus, classification of invariant H -solutions is reduced to the problem of classification of subgroups of G , up to similarity. An optimal system of s -dimensional subgroups of G is a list of conjugacy inequivalent s -dimensional subgroups of G with the property that any other subgroup is conjugate to precisely one subgroup in the list. Similarly, a list of s -dimensional subalgebras forms an optimal system if every s -dimensional subalgebra of $\mathfrak{g}$ is equivalent to a unique member of the list under some element of the adjoint representation: $\tilde{\mathfrak{h}} = \text{Ad}(g) \cdot \mathfrak{h}$.

Let H and $\tilde{H}$ be connected, s -dimensional Lie subgroups of the Lie group G with corresponding Lie subalgebras $\mathfrak{h}$ and $\tilde{\mathfrak{h}}$ of the Lie algebra $\mathfrak{g}$ of G . Then $\tilde{H} = gHg^{-1}$ are conjugate subgroups if and only if $\tilde{\mathfrak{h}} = \text{Ad}(g) \cdot \mathfrak{h}$ are conjugate subalgebras (Proposition 3.7 of [4]). Thus, the problem of finding an optimal system of subgroups is equivalent to that of finding an optimal system of subalgebras, and so we concentrate on it.

For one-dimensional subalgebras, the classification problem is essentially the same as the problem of classifying the orbits of the adjoint representation, since each one-dimensional subalgebra is determined by a nonzero vector in $\mathfrak{se}(3)$ and so to "simplify" it as much as possible.

The adjoint action is given by the Lie series

$$\text{Ad}(\exp(s.X_i).X_j) = X_j - s.[X_i, X_j] + \frac{s^2}{2}.[X_i, [X_i, X_j]] - \dots, \quad (10)$$

where $[X_i, X_j]$ is the commutator for the Lie algebra, s is a parameter, and $i, j = 1, \dots, 6$ ([4], page 199). We can write the adjoint action for the Lie algebra $\mathfrak{se}(3)$, and show that

3.0.0.4 Theorem *A one-dimensional optimal system of $\mathfrak{se}(3)$ is given by*

$$\begin{aligned} A_1^1 : X_6, & \quad A_1^3 : X_2 + b X_5, & A_1^5 : X_1 + a X_2 + b X_6, & \quad A_1^7 : X_2 + a X_3 + b X_5, \\ A_1^2 : X_1 + b X_4, & A_1^4 : X_3 + b X_6, & A_1^6 : X_1 + a X_3 + b X_4, \end{aligned} \quad (11)$$

where $a, b \in \mathbb{R}$ and $a \neq 0$.

Proof: $F_i^s : \mathfrak{se}(3) \rightarrow \mathfrak{se}(3)$ defined by $X \mapsto \text{Ad}(\exp(s_i X_i).X)$ is a linear map, for $i = 1, \dots, 6$. The matrix M_i^s of F_i^s , $i = 1, \dots, 6$, with respect to basis $\{X_1, \dots, X_6\}$ is

$$\begin{aligned} M_1^s &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & s & 0 & 1 & 0 \\ 0 & -s & 0 & 0 & 0 & 1 \end{bmatrix}, & M_2^s &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -s & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ s & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, & M_3^s &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & s & 0 & 1 & 0 & 0 \\ -s & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \\ M_4^s &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & C & S & 0 & 0 & 0 \\ 0 & S & C & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & C & S \\ 0 & 0 & 0 & 0 & -S & C \end{bmatrix}, & M_5^s &= \begin{bmatrix} C & 0 & -S & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ S & 0 & C & 0 & 0 & 0 \\ 0 & 0 & 0 & C & 0 & -S \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & S & 0 & C \end{bmatrix}, & M_6^s &= \begin{bmatrix} C & S & 0 & 0 & 0 & 0 \\ -S & C & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & C & S & 0 \\ 0 & 0 & 0 & -S & C & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}. \end{aligned} \quad (12)$$

respectively, where $S = \sin s$ and $C = \cos s$. Let $X = \sum_{i=1}^6 a_i X_i$, then

$$F_6^{s_6} \circ \dots \circ F_1^{s_1} : X \mapsto (\cos s_5 \cdot \cos s_6 \cdot a_1 + \cos s_5 \cdot \sin s_6 \cdot a_2 - \sin s_5 \cdot a_3) \cdot X_1 + \dots \quad (13)$$

Now, we can simplify X as follows:

If a_1, a_2 and $a_3 = 0$, then we can make the coefficients of $X_1, \dots, X_5$ vanish, by $s_4 = -\arctan(a_5/a_6)$ and $s_5 = \arctan(a_4/a_6)$. Scaling X if necessary, we can assume that $a_6 = 1$. And X is reduced to the Case of A_1^1 .

If a_2 and $a_3 = 0$ but $a_1 \neq 0$, then we can make the coefficients of X_2, X_3, X_5 and X_6 vanish, by $s_2 = -a_6/a_1$ and $s_3 = a_5/a_1$. Scaling X if necessary, we can assume that $a_1 = 1$. And X is reduced to the Case of A_1^2 .

If a_1 and $a_3 = 0$ but $a_2 \neq 0$, then we can make the coefficients of X_1, X_2, X_3 , and X_6 vanish, by $s_1 = a_6/a_2$ and $s_3 = -a_4/a_2$. Scaling X if necessary, we can assume that $a_3 = 1$. And X is reduced to the Case of A_1^3 .

If a_1 and $a_2 = 0$ but $a_3 \neq 0$, then we can make the coefficients of X_1, X_2, X_4 and X_5 vanish, by $s_1 = -a_5/a_3$ and $s_2 = a_4/a_3$. Scaling X if necessary, we can assume that $a_3 = 1$. And X is reduced to the Case of A_1^4 .

If a_1 and $a_2 \neq 0$, then we can make the coefficients of X_3 , X_4 and X_5 vanish, by $s_1 = a_6/a_2$, $s_3 = a_5/a_1$ and $s_4 = -\arctan(a_3/a_2)$. Scaling X if necessary, we can assume that $a_1 = 1$. And X is reduced to the Case of A_1^5 .

If $a_2 = 0$ but a_1 and $a_3 \neq 0$, then we can make the coefficients of X_2 , X_5 and X_6 vanish, by $s_1 = -a_5/a_3$, $s_2 = -a_6/a_1$ and $s_4 = -\arctan(a_3/a_2)$. Scaling X if necessary, we can assume that $a_1 = 1$. And X is reduced to the Case of A_1^6 .

If $a_1 = 0$ but a_2 and $a_3 \neq 0$, then we can make the coefficients of X_3 , X_4 and X_5 vanish, by $s_1 = a_6/a_2$, and $s_2 = a_4/a_3$. Scaling X if necessary, we can assume that $a_2 = 1$. And X is reduced to the Case of A_1^7 . $\square$

3.0.0.5 Theorem A two-dimensional optimal system of $\mathfrak{se}(3)$ is given by

$$\begin{aligned} A_2^1 : X_2, X_5, & \quad A_2^3 : X_1, X_2 + a X_4, & \quad A_2^5 : X_2, X_3 + a X_5, \\ A_2^2 : X_3, X_6, & \quad A_2^4 : X_1, X_3 + a X_4, & \quad A_2^6 : X_3, X_1 + a X_4, \end{aligned} \quad (14)$$

where $a \in \mathbb{R}$. All of these sub-algebras are Abelian.

Proof: In this proof, we shall assume that $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X, Y\}$ is a 2-dimensional Lie sub-algebra of $\mathfrak{se}(3)$.

Let $X = X_6$ be as the Case of A_1^1 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_3 X_3 + b_6 X_6$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $Y = X_3$. Now $\mathfrak{g}$ is reduced to the Case of A_2^2 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$ defined as (12).

Let $X = X_1 + a_6 X_6$ be as the Case of A_1^2 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_3 X_3 + b_1 X$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $Y = X_3$. Now $\mathfrak{g}$ is reduced to the Case of A_2^2 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$.

Let $X = X_2 + a_5 X_5$ be as the Case of A_1^3 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_2 X_2 + b_5 X_5$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_2$ and $Y = X_5$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^1 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$.

Let $X = X_3 + a_6 X_6$ be as the Case of A_1^4 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_3 X_3 + b_6 X_6$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_6$ and $Y = X_3$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^2 .

Let $X = X_1 + a_2 X_2 + a_4 X_4$ be as the Case of A_1^5 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_1 X_1 + (b_2/a_2) X$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_1$ and $Y = X_2 + a X_4$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^3 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$.

Let $X = X_2 + a_3 X_3 + a_4 X_4$ be as the Case of A_1^6 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_1 X_1 + (b_3/a_3) X$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_1$ and $Y = X_3 + a X_4$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^4 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$.

Let $X = X_2 + a_3 X_3 + a_5 X_5$ be as the Case of A_1^7 , $Y = \sum_{i=1}^6 b_i X_i$, and $[X, Y] = \alpha X + \beta Y$. Then, we have $Y = b_2 X_2 + (b_3/a_3) X$ and $\alpha = \beta = 0$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_2$ and $Y = X_3 + a X_5$. Now, $\mathfrak{g}$ is reduced to the Case of A_2^5 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$. And, we have proved the Theorem. $\square$

3.0.0.6 Theorem A three-dimensional optimal system of $\mathfrak{se}(3)$ is given by $A_3 : X_1 + a X_4, X_2, X_3$, where $a \in \mathbb{R}$. The commutator table of A_3 is Table 2.

Table 2
Commutation table of A_3 .

	X	Y	Z
X	0	$-a$	0
Y	a	0	0
Z	0	0	0

Table 3
Commutation table of A_4 .

	X_1	X_2	X_3	X_4
X_1	0	0	0	0
X_2	0	0	0	X_3
X_3	0	0	0	$-X_2$
X_4	0	$-X_3$	X_2	0

Proof: In this proof, we assume that $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X, Y, Z\}$ is a 3-dimensional Lie sub-algebra of $\mathfrak{se}(3)$.

Let $X = X_2$ and $Y = X_5$ are as the Case of A_2^1 , $Z = \sum_{i=1}^6 b_i X_i$, $[X, Z] = \alpha_1 X + \beta_2 Y + \beta_3 Z$ and $[Y, Z] = \beta_1 X + \beta_2 Y + \beta_3 Z$. Then, we have $Z = b_2 X + b_5 Y$. By a suitable change of base of $\mathfrak{g}$, we can assume that $Z = 0$, and $\mathfrak{g}$ is not a 3-dimensional sub-algebra. Thus, in this case we have not any 3-dimensional sub-algebra.

The Cases A_2^i , $i = 2, 3, 4, 5$ are similar. Thus, in these cases we have not any 3-dimensional sub-algebra.

Let $X = X_3$ and $Y = X_1 + a X_4$ are as the Case of A_2^6 , $Z = \sum_{i=1}^6 b_i X_i$, $[X, Z] = \alpha_1 X + \beta_2 Y + \beta_3 Z$ and $[Y, Z] = \beta_1 X + \beta_2 Y + \beta_3 Z$. Then, we have $Z = (b_1/\alpha_3) \cdot (-\beta_3 X + \alpha_3 Y - a X_2)$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_1 + a X_4$, $Y = X_2$ and $Z = X_3$. Now, $\mathfrak{g}$ is reduced to the Case of A_3 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$ defined as (12). $\square$

3.0.0.7 Theorem A four-dimensional optimal system of $\mathfrak{se}(3)$ is given by $A_4 : X_1, X_2, X_3, X_4$. The commutator table of A_4 is Table 3. *Proof:* Assume that $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X_1 + a X_4, X_2, X_3, X\}$ be a Lie sub-algebra of $\mathfrak{se}(3)$, where

$X = \sum_{i=1}^6 b_i X_i$. Then, we have $X = \sum_{i=1}^4 b_i X_i$. By a suitable change of base of $\mathfrak{g}$, we can assume that $X = X_4$. Now, $\mathfrak{g}$ is reduced to the Case of A_4 . We can not be used to further simplify this sub-algebra, by $F_i^{s_i}$, $i = 1, \dots, 6$ defined as (12). $\square$

3.0.0.8 Theorem $\mathfrak{se}(3)$ has not any five-dimensional Lie sub-algebra.

Proof: Assume that $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X_1, X_2, X_3, X_4, X\}$ be a Lie sub-algebra of $\mathfrak{se}(3)$, where $X = \sum_{i=1}^6 b_i X_i$. Then, we have $X = \sum_{i=1}^4 b_i X_i$, and $\mathfrak{g} = \text{Span}_{\mathbb{R}}\{X_1, X_2, X_3, X_4\}$ is not a 5-dimensional sub-algebra. $\square$

4 Invariant solutions

To obtain the group transformation which is generated by the infinitesimal generators $X_i = \sum_{j=1}^3 \xi_j^i + \phi^i \partial_u$ for $i = 1, \dots, 6$, we need to solve the 6 systems of first order ordinary differential equations,

$$\begin{aligned} \tilde{x}'_j(s) &= \xi_j^i(\tilde{x}_1(s), \tilde{x}_2(s), \tilde{x}_3(s), \tilde{u}(s)), \quad \tilde{x}_j(0) = x_j, \quad i = 1, \dots, 6 \\ \tilde{u}'(s) &= \phi^i(\tilde{x}_1(s), \tilde{x}_2(s), \tilde{x}_3(s), \tilde{u}(s)), \quad \tilde{u}(0) = u. \quad j = 1, 2, 3 \end{aligned} \tag{15}$$

Exponentiating the infinitesimal symmetries of the Poisson's equation (1), we get the one parameter groups $g_k(s)$ generated by X_k for $k = 1, \dots, 6$; and consequently, we have

4.0.0.9 Theorem If $u = h(x, y, z)$ is a solution of the Poisson's equation (1), so are the functions

$$\begin{aligned} g_1(s) \cdot h(x, y, z) &= h(x + s, y, z), & g_4(s) \cdot h(x, y, z) &= h(x, y \cos s - z \sin s, z \cos s + y \sin s), \\ g_2(s) \cdot h(x, y, z) &= h(x, y + s, z), & g_5(s) \cdot h(x, y, z) &= h(x \cos s + z \sin s, y, z \cos s - x \sin s), \\ g_3(s) \cdot h(x, y, z) &= h(x, y, z + s), & g_6(s) \cdot h(x, y, z) &= h(x \cos s - y \sin s, x \sin s + y \cos s, z), \end{aligned} \quad (16)$$

References

- [1] G.W. BLUMAN, J.D. COLE, *Similarity Methods for Differential Equations*, Springer, Berlin, 1974.
- [2] J. JOST, *Partial differential equations*, Springer, GTM 214, 2002.
- [3] M. NADJAFIKHAH, *Lie Symmetries of Inviscid Burgers Equation*, Adv. appl. Clifford alg., DOI 10.1007/s00006-003-0000.
- [4] P.J. OLVER, *Applications of Lie Groups to Differential Equations*, Springer, New York, 1986.
- [5] L.V. OVSIANNIKOV, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.
- [6] A.D. POLYANIN, and V.F. ZAITSEV, *Handbook of Nonlinear Partial Differential Equations*, Chapman & Hall/CRC, Boca Raton, 2004.
- [7] R. SESHADRI and T.Y. NA, *Group Invariance in Engineering Boundary Value Problems*, Springer-Verlag, New York, 1985.
- [8] T.R. SEKHAR, V.D. SHARMA, *Similarity solutions for three dimensional Euler equations using Lie group analysis*, Appl. Math. and Comp. 196 (2008) 147–157.